

Gravitational Zero Point Energy induces Physical Observables

REMO GARATTINI

*Università degli Studi di Bergamo, Facoltà di Ingegneria,
Viale Marconi 5, 24044 Dalmine (Bergamo) ITALY.
INFN - sezione di Milano, Via Celoria 16, Milan, Italy
remo.garattini@unibg.it*

We consider the contribution of Zero Point Energy on the induced Cosmological Constant and on the induced Electric/Magnetic charge in absence of matter fields. The method is applicable to every spherically symmetric background. Extensions to a generic $f(R)$ theory are also allowed. Only the graviton appears to be fundamental to the determination of Zero Point Energy.

The path integral approach to quantum gravity

$$Z = \int \mathcal{D}[g_{\mu\nu}] \exp iS_g[g_{\mu\nu}] \quad (1)$$

is a powerful method to study the quantization of the gravitational field, especially in the context of a WKB approximation on a given background. Indeed, if one considers a background $\bar{g}_{\mu\nu}$, the gravitational field splits into $g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$, where $h_{\mu\nu}$ is a quantum fluctuation around the background field and Eq. (1) becomes

$$\int \mathcal{D}g_{\mu\nu} \exp iS_g[g_{\mu\nu}] \simeq \exp iS_g[\bar{g}_{\mu\nu}] \int \mathcal{D}h_{\mu\nu} \exp iS_g^{(2)}[h_{\mu\nu}], \quad (2)$$

where $S_g^{(2)}[h_{\mu\nu}]$ is the action approximated to second order. Since the second order action is quadratic in $h_{\mu\nu}$, the integration in Eq.(2) is straightforward. In absence of matter fields, one may wonder what is the contribution of quantum fluctuations to the cosmological constant, by computing the averaged energy-momentum tensor

$$\langle T_{\mu\nu}^{C.C.} \rangle = -\frac{2}{\sqrt{-g}} \frac{\delta \ln Z}{\delta g^{\mu\nu}}, \quad (3)$$

where $T_{\mu\nu}^{C.C.}$ is the energy momentum tensor related to the cosmological term. A general solution of Eq.(3) is a complicated task, however the Arnowitt-Deser-Misner ($\mathcal{ADM}$) variables offer a valid example to extract the relevant degrees of freedom.¹ In terms of these variables, the metric background is

$$ds^2 = -N^2 dt^2 + g_{ij} (N^i dt + dx^i) (N^j dt + dx^j) = -N^2 dt^2 + \frac{dr^2}{1 - \frac{b(r)}{r}} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (4)$$

We recognize that the *lapse function* N is invariant and the *shift function* N_i is absent. To have an effective reduction of the modes, we consider perturbations of the gravitational field on the hypersurface $\Sigma \subset \mathcal{M}$. This means that we are “freezing” the perturbation of the lapse and the shift functions respectively. In summary,

$$\begin{cases} g_{ij} \longrightarrow \bar{g}_{ij} + h_{ij} \\ N \longrightarrow N \\ N_i \longrightarrow 0 \end{cases}. \quad (5)$$

If we indicate with u_μ a time-like unit vector, then it is possible to prove that²

$$\langle T_{\mu\nu}^{C.C.} \rangle = -\frac{i}{2} g_{\mu\nu} \int \frac{d^4 k}{(2\pi)^4} \ln \lambda_{TT}^2 + \frac{i}{2} (g_{\mu\nu} + u_\mu u_\nu) \int \frac{d^3 k}{(2\pi)^3} \ln \lambda_{V^\perp}^2, \quad (6)$$

where λ_{TT} are the transverse-traceless (TT) eigenvalues of the following second order differential operator

$$O^{ijkl} h_{kl} = \left[\Delta_L^{ijkl} + 4R^{ij} g^{kl} + \frac{1}{N^2} \frac{\partial^2}{\partial t^2} g^{ik} g^{jl} \right] h_{kl} = -(\lambda_{TT}^2) h^{ij} \quad (7)$$

and Δ_L is the Lichnerowicz operator. From Eq.(6), integrating out the time component, the energy density simply becomes

$$\frac{\Lambda}{8\pi G} = -\frac{i}{2} \left[\int \frac{d^4 k}{(2\pi)^4} \ln \lambda_{TT}^2 \right] = -\frac{1}{2} \sum_{i=1}^2 \int \frac{d^3 k}{(2\pi)^3} \sqrt{\lambda_i^2(|\vec{k}|)}. \quad (8)$$

The same expression can be obtained by computing

$$\frac{1}{V} \frac{\int \mathcal{D}[g_{ij}] \Psi^*[g_{ij}] \int_\Sigma d^3 x \mathcal{H} \Psi[g_{ij}]}{\int \mathcal{D}[g_{ij}] \Psi^*[g_{ij}] \Psi[g_{ij}]} = \frac{1}{V} \frac{\langle \Psi | \int_\Sigma d^3 x \hat{\Lambda}_\Sigma | \Psi \rangle}{\langle \Psi | \Psi \rangle} = -\frac{\Lambda}{\kappa}, \quad (9)$$

where we have integrated over a hypersurface Σ , divided by its volume and functionally integrated over quantum fluctuation with the help of some trial wave functionals. $\mathcal{H}$ is the hamiltonian constraint without the cosmological term. Note that Eq.(9) can be derived starting with the Wheeler-De Witt equation (WDW)³ which represents invariance under *time* reparametrization. Eq.(9) represents the Sturm-Liouville problem associated to the cosmological constant. The related boundary conditions are dictated by the choice of the trial wavefunctionals which, in our case are of the Gaussian type. Different types of wavefunctionals correspond to different boundary conditions. The evaluation of the r.h.s. of Eq.(8) is performed via the absorption of the divergent part into the re-definition of the bare classical constant $\Lambda \rightarrow \Lambda_0 + \Lambda^{div}$. The dependence on the arbitrary mass scale μ can be removed with the help of a renormalization group-like equation. Solving it we find that the renormalized constant Λ_0 should be treated as a running one in the sense that it varies provided that the scale μ is changing. Finally, we obtain

$$\frac{\Lambda_0(\mu_0, r)}{8\pi G} = -\frac{1}{64\pi^2} \sum_{i=1}^2 \left[m_i^4(r) \ln \left(\frac{m_i^2(r)}{4\mu_0^2} \sqrt{e} \right) \right], \quad (10)$$

where $m_i^2(r)$ play the rôle of an effective mass with

$$\begin{cases} m_1^2(r) = \frac{6}{r^2} \left(1 - \frac{b(r)}{r} \right) + \left[\frac{3}{2r^2} b'(r) - \frac{3}{2r^3} b(r) \right] \\ m_2^2(r) = \frac{6}{r^2} \left(1 - \frac{b(r)}{r} \right) + \left[\frac{1}{2r^2} b'(r) + \frac{3}{2r^3} b(r) \right] \end{cases}. \quad (11)$$

Thus, by changing the form of $b(r)$, one can explore the spectrum of $\Lambda_0(\mu_0, r)/8\pi G$. For example, we find:^{4,5}

Schwarzschild/Naked	$b_S(r) = 2MG$	$b_{Naked-S}(r) = -2MG$
dS/AdS	$b_{dS}(r) = \frac{\Lambda_{dS}}{3}r^3$	$b_{AdS}(r) = -\frac{\Lambda_{AdS}}{3}r^3$
SdS/S-AdS	$b_{SdS}(r) = 2MG + \frac{\Lambda_{dS}}{3}r^3$	$b_{SAdS}(r) = 2MG - \frac{\Lambda_{AdS}}{3}r^3$

Note that the method represented by Eqs.(3,9) can be applied even to the electric/magnetic charge by replacing in Eq.(10) the induced cosmological constant with the induced electric/magnetic charge. In doing so we obtain the quantum gravitational contribution to such observables⁶

$$Q_{e,0}^2(\mu_0, r) = -\frac{r^4}{32\pi} \sum_{i=1}^2 \left[m_i^4(r) \ln \left(\frac{m_i^2(r)}{4\mu_0^2} \sqrt{e} \right) \right]. \quad (12)$$

Note the difference between Eq.(10) and Eq.(12). In the latter the presence of the factor r^4 at the numerator allows a less singular behavior close to the throat. A bound on $Q_{e,0}^2(\mu_0, r)$ shows that

$$2.2 \times 10^{-2} = \frac{9}{128\pi} \geq Q_{e,0}^2(\mu_0, \bar{r}) \geq \frac{9}{200\pi} = 1.4 \times 10^{-2}. \quad (13)$$

Note that the fine structure constant is $\frac{1}{137} = .73 \times 10^{-2}$. Note also that Zero Point Energy allows an “*induced magnetic charge*”, at least in principle. It is interesting that in all these examples except the Schwarzschild case one could impose a running of the Newton constant G instead of $\Lambda_0(\mu_0, r)/8\pi G$.⁵ An extension of the method which considers a generic function of the curvature $f(R)$ has been discussed in Refs.^{5,7} The modified gravity model induces a modification on $\Lambda_0(\mu_0, r)/8\pi G$ which is strongly dependent on the kind of theory considered. How this affects the induced Electric/Magnetic charge is an open problem.

References

1. R. Arnowitt, S. Deser, and C. W. Misner, in *Gravitation: An Introduction to Current Research*, edited by L. Witten (John Wiley & Sons, Inc., New York, 1962). ArXiv:gr-qc/0405109.
2. R. Garattini, *Wormholes or Gravastars?* arXiv:1001.3831 [gr-qc].
3. B. S. DeWitt, *Phys. Rev.* **160**, 1113 (1967).
4. R. Garattini, *TSPU Vestnik* **44 N7**, 72 (2004); gr-qc/0409016. R. Garattini, *J. Phys. Conf. Ser.* **33** 215 (2006); gr-qc/0510062. R. Garattini, *J. Phys. A* **39** 6393, (2006); gr-qc/0510061. R. Garattini, *The induced Cosmological Constant as a tool for exploring geometries*. arXiv:0812.4037 [gr-qc].
5. R. Garattini, *Running Cosmological Constant and Running Newton Constant in Modified Gravity Theories*. arXiv:0911.2393 [gr-qc].
6. R. Garattini, *Phys. Lett. B* **666** 189 (2008). arXiv:0807.0082 [gr-qc].
7. S. Capozziello and R. Garattini, *Class.Quant.Grav.* **24**, 1627 (2007); gr-qc/0702075. R. Garattini, *J.Phys.Conf.Ser.*174 012066 (2009); arXiv:0902.1087 [gr-qc].